arXiv:1202.5457v1 [math.CA] 14 Feb 2012

On the Equivalence of Fourier Expansion and Poisson Summation Formula for the Series Approximation of the Exponential Function

S. M. Abrarov[*], **B. M. Quine**[*][†]**and R. K. Jagpal**[†]

February 14, 2012

Abstract

In this short note we show the equivalence of Fourier expansion and Poisson summation approaches for the series approximation of the exponential function $\exp\left(-t^2/4\right)$. The application of the Poisson summation formula is shown to reduce to that of the Fourier expansion method.

Keywords: Fourier expansion, Poisson summation formula, Exponential function

1 Methodology

Let us apply the Poisson summation formula (see for example [1]) to the exponential function $\exp\left(-t^2/4\right) \equiv \exp\left[-(t/2)^2\right]$, i.e.:

$$\sum_{n=-\infty}^{\infty} \exp\left[-\left(\frac{t}{2}+n\tau_m\right)^2\right] = \frac{\sqrt{\pi}}{\tau_m}\sum_{n=-\infty}^{\infty} \exp\left(-\frac{\pi^2 n^2}{\tau_m^2}\right)\cos\left(\frac{\pi n}{\tau_m}t\right)$$

---

[*] Dept. Earth and Space Science and Engineering, York University, Toronto, Canada, M3J 1P3.

[†] Dept. Physics and Astronomy, York University, Toronto, Canada, M3J 1P3.

or

$$\begin{aligned}\exp\left(-\frac{t^2}{4}\right) = {} & \frac{\sqrt{\pi}}{\tau_m}\sum_{n=-\infty}^{\infty}\exp\left(-\frac{\pi^2 n^2}{\tau_m^2}\right)\cos\left(\frac{\pi n}{\tau_m}t\right) \\ & -\left\{\sum_{n=-\infty}^{-1}\exp\left[-\left(\frac{t}{2}+n\tau_m\right)^2\right]+\sum_{n=1}^{\infty}\exp\left[-\left(\frac{t}{2}+n\tau_m\right)^2\right]\right\},\end{aligned} \tag{1}$$

where $2\tau_m$ is the period. Assuming that the half-period is large enough, say $\tau_m \geqslant 12$, within the range $t \in [-\tau_m, \tau_m]$ it follows that

$$\exp\left(-\frac{t^2}{4}\right) >> \exp\left[-\left(\frac{t}{2}\pm\tau_m\right)^2\right] >> \exp\left[-\left(\frac{t}{2}\pm 2\tau_m\right)^2\right] >> \exp\left[-\left(\frac{t}{2}\pm 3\tau_m\right)^2\right] \ldots .$$

Consequently, we can ignore the terms in curly brackets in equation (1) and approximate exponential function as

$$\exp\left(-\frac{t^2}{4}\right) \approx \frac{\sqrt{\pi}}{\tau_m}\sum_{n=-\infty}^{\infty}\exp\left(-\frac{\pi^2 n^2}{\tau_m^2}\right)\cos\left(\frac{\pi n}{\tau_m}t\right). \tag{2}$$

Taking into account that

$$\exp\left(-\frac{\pi^2(-n)^2}{\tau_m^2}\right)\cos\left(\frac{\pi\,(-n)}{\tau_m}t\right) \equiv \exp\left(-\frac{\pi^2 n^2}{\tau_m^2}\right)\cos\left(\frac{\pi n}{\tau_m}t\right),$$

the approximation (2) can be rearranged in form

$$\exp\left(-\frac{t^2}{4}\right) \approx \frac{\sqrt{\pi}}{\tau_m} + \frac{2\sqrt{\pi}}{\tau_m}\sum_{n=1}^{\infty}\exp\left(-\frac{\pi^2 n^2}{\tau_m^2}\right)\cos\left(\frac{\pi n}{\tau_m}t\right)$$

or

$$\exp\left(-\frac{t^2}{4}\right) \approx -\frac{\sqrt{\pi}}{\tau_m} + \frac{2\sqrt{\pi}}{\tau_m}\sum_{n=0}^{\infty}\exp\left(-\frac{\pi^2 n^2}{\tau_m^2}\right)\cos\left(\frac{\pi n}{\tau_m}t\right).$$

Lastly, defining

$$a_n = \frac{2\sqrt{\pi}}{\tau_m}\exp\left(-\frac{\pi^2 n^2}{\tau_m^2}\right)$$

we obtain

$$\exp\left(-\frac{t^2}{4}\right) \approx -\frac{a_0}{2} + \sum_{n=0}^{\infty} a_n \cos\left(\frac{\pi n}{\tau_m}t\right), \qquad t \in [-\tau_m, \tau_m].$$

This approximation is proposed recently for efficient computation of the Voigt function [2].

2 Conclusion

We show the equivalence of series approximation derived by Fourier expansion and by Poisson summation formula for the exponential function $\exp\left(-t^2/4\right)$.

References

[1] R. J. Beerends, H. G. ter Morsche, J. C. van den Berg and E. M. van de Vrie, Fourier and Laplace Transforms, Cambridge University Press, 2003.

[2] S. M. Abrarov, B. M. Quine and R. K. Jagpal, Rapidly convergent series for high-accuracy calculation of the Voigt function, J. Quant. Spect. Rad. Trans., 111 (2010) 372-375.